

# (12) UK Patent Application (19) GB (11) 2 317 173 (13) A

(43) Date of A Publication 18.03.1998

(21) Application No 9719084.7

(22) Date of Filing 09.09.1997

(30) Priority Data

(31) 60025927

(32) 12.09.1996

(33) US

(71) Applicant(s)

Merck & Co Inc

(Incorporated in USA - New Jersey)

P O Box 2000, 126 East Lincoln Avenue, Rahway,  
New Jersey 07065-0900, United States of America

(72) Inventor(s)

Ronald M Kim

(74) Agent and/or Address for Service

I J Hiscock

Merck & Co. Inc, European Patent Department,  
Terlings Park, Eastwick Road, HARLOW, Essex,  
CM20 2QR, United Kingdom

(51) INT CL<sup>6</sup>

C07C 43/115 61/00

(52) UK CL (Edition P)

C2C CAA CWR C22Y C220 C226 C227 C30Y C348 C36Y  
C360 C362 C364 C403 C406 C43X C496 C50Y C500  
C643 C644 C652 C662 C699 C778 C779

(56) Documents Cited

None


(58) Field of Search

UK CL (Edition O) C2C CWR  
INT CL<sup>6</sup> C07B 61/00, C07C 43/00  
ONLINE: CAS- ONLINE, EDOC, JAPIO, WPI

## (54) POLYARYL-POLY (ETHYLENE GLYCOL) SUPPORTS FOR SOLUTION-PHASE COMBINATORIAL SYNTHESIS


(57) The present invention is directed to the certain polyaryl-poly(ethylene glycol) compounds ("polyaryl-PEG" compounds) which are useful as soluble polyvalent supports in the preparation of combinatorial libraries of compounds.

The compounds of Formula I, II or III:



wherein:

A, A', A'' and A''' are independently selected from:



each X and X' are -OH, -Br, -Cl, -I, -SH, -CO<sub>2</sub>H, -CO<sub>2</sub>Cl, CO<sub>2</sub>Br, -SO<sub>3</sub>H, -SO<sub>2</sub>Cl, -NH<sub>2</sub>, and -CH=CH<sub>2</sub>;

each Y is hydrogen, -CH<sub>3</sub>, -O-(C<sub>1-6</sub>alkyl) and -O(CH<sub>2</sub>CH<sub>2</sub>O)<sub>n</sub>-CH<sub>3</sub>;

j and j' are each 0 to 4;

k and k' are each 1 to 12;

m and m' are each 1 to 4; and

each n is 0 to 12; and salts thereof.

GB 2 317 173 A

## TITLE OF THE INVENTION

POLYARYL-POLY(ETHYLENE GLYCOL) SUPPORTS FOR  
SOLUTION-PHASE COMBINATORIAL SYNTHESIS

## 5 CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority under 35 U.S.C. § 119(e)  
from USSN 60/025,927, filed September 12, 1996.

## BACKGROUND OF THE INVENTION

10 The standard method for searching for new chemical  
compounds which can effectively modulate biological processes  
employs the screening of pre-existing compounds in assays which  
have been designed to test particular properties of the compound being  
15 screened. Similarly, in designing compounds having desired  
physiochemical properties for general chemical applications,  
numerous compounds must be individually prepared and tested.

To reduce the time and expense involved in preparing and  
screening a large number of compounds for biological activity or for  
desirable physiochemical properties, technology has been developed  
20 for providing libraries of compounds for the discovery of lead  
compounds. Current methods for generating large numbers of  
molecularly diverse compounds focus on the use of solid phase  
synthesis. The generation of combinatorial libraries of chemical  
compounds by employing solid phase synthesis is well known in the  
25 art. For example, Geysen, et al. (Proc. Natl. Acad. Sci. USA, 3998  
(1984) describe the construction of multi-amino acid peptide libraries;  
Houghton, et al. (Nature, 354, 84 (1991) and PCT Patent Pub. No. WO  
92/09300) describe the generation and use of synthetic peptide  
combinatorial libraries for basic research and drug discovery; Lam, et  
30 al. (Nature, 354, 82 (1991) and PCT Patent Pub. No. WO 92/00091)  
describe a method of synthesis of linear peptides on a solid support  
such as polystyrene or polyacrylamide resin.

The growing importance of combinatorial chemistry as an  
integral component of the drug discovery process has spurred extensive

technological and synthetic advances in the field (Thompson, L. A.; Ellman, J. A. (1996) *Chem. Rev.* 96, 555-600). Founded in peptide synthesis devised by Merrifield, solid phase chemistry has emerged as the preeminent method for construction of small molecule combinatorial

5 libraries (see e.g. Merrifield, R. B. (1963) *J. Am. Chem. Soc.* 85, 2149-2154; (a) Terrett, N. K.; Gardner, M.; Gordon, D. W.; Kobylecki, R. J.; Steele, J. (1995) *Tetrahedron* 51(30), 8135-8173. (b) Gordon, E. M.; Barrett, R. W.; Dower, W. J.; Fodor, S. P. A.; Gallop, M. A. (1994) *J. Med. Chem.* 37, 1385-1401.).

10 It is known that a wide variety of organic reactions can be carried out on substrates immobilized on resins. These include, in addition to peptide synthesis, nucleophilic displacements on benzylic halides, halogenation, nitration, sulfonation, oxidation, hydrolysis, acid chloride formation, Friedel-Crafts reactions, reductions, metallation, and

15 the like which are well known in the art. (See for example, Mathur, et al., "Polymers as Aids in Organic Chemistry", Academic Press, New York, 18 (1980) and Farrall, et al., *J. Org. Chem.*, 41, 3877 (1976)).

One variant on the use of substrates immobilized on resin beads employs polystyrene pins as supports for the substrates (see

20 Geysen, et al., *J. Immunol. Meth.*, 102, 259 (1987) and another variant employs a flat solid support in the form of a tape or a streamer (see PCT Patent Publication WO 96/16078, published May 30, 1996).

Although combinatorial chemistry may be attempted by employing traditional synthetic chemistry in the solution phase, this is

25 essentially impractical due to the difficulty in separating complex mixtures of intermediates and final compounds from reagents and solvents. Traditional solution phase chemistry has been criticized as being unsuitable as a technique which aims to simultaneously produce a multiplicity of new products, since this would not allow physical

30 separation among the different products. The products are therefore likely to be contaminated with excess reagents, by-products, etc., leading to difficulties in separation and purification.

Central to the power of solid phase synthesis is the ease by which reagents and solvents are removed simply by washing. This allows

for the purification of resin-bound mixtures of great complexity, and the use of large reagent excesses to drive reactions to completion. The "infinite dilution" obtained on solid supports can also prevent side reactions which may occur in solution. Despite its advantages, nontrivial liabilities are associated with solid phase synthesis. Most notable is the often arduous task of modifying solution phase chemistry to the solid phase, with its potential pitfalls such as poor solvation, differential site accessibility, and incompatibility of the polymer support with reagents or reaction conditions. Often the most time-consuming aspect of combinatorial library synthesis is not construction of the library itself, but rather translation of solution phase chemistry to the solid phase. Although a significant portion of organic chemistry can be adapted to synthesis on the solid phase, some reactions, such as heterogeneous catalysis, would be exceedingly difficult to conduct on reagents linked to a solid phase support. In addition to synthetic complications which may arise from employing a solid support, few analytical techniques exist for characterization of resin-bound compounds. Even if NMR or IR analytical methods are attempted, they usually require specialized equipment and techniques and often yield spectra of low quality. Thus, it is very difficult to monitor reactions conducted on solid phase. Even analysis of cleaved intermediates can be ambiguous, since the harsh cleavage conditions that are often required may be detrimental to the molecules of interest.

Accordingly, alternative methodology for the preparation of combinatorial libraries which overcomes the drawbacks of solid phase synthesis would provide a significant advance in the field.

Classical solution phase techniques have been developed in an attempt to overcome the drawbacks of solid phase synthesis. One approach is to separate acidic and basic reagents from the resultant products by adding a water solution of additional acids or bases (See Boger, et al, *J. Am. Chem. Soc.*, (February 28, 1996)). A variant of this method takes advantage of extraction of the reagents or products into a perfluorocarbon solvent.

Another approach, termed liquid-phase combinatorial synthesis (LPCS), in which combinatorial libraries are synthesized on soluble polyethylene glycol (PEG) supports has been recently described (Han, H.; Wolfe, M.M.; Brenner, S.; Janda, K.D. *Proc. Natl. Acad. Sci. USA*, 92, 6419-6423 (1995) and Han, H.; Janda, K.D. *J. Am. Chem. Soc.*, 118, 2539-2544 (1996)). In this method, monofunctional PEG which falls within a certain molecular weight range is used as a support for synthetic reactions. When the reactions are complete, ether is added to the solution which causes precipitation of the PEG, which is then isolated by filtration. This precipitation/ crystallization of the PEG-protected molecules from ether allows for removal of reagents and solvents by filtration, thus combining the advantages of solution phase chemistry and the utility of solid phase purification. This technology has been reviewed in *Science*, 272, 1266-1268 (May 31, 1996).

Certain polyvalent poly(ethylene glycol) compounds have been disclosed, but they have not been previously noted as being useful as soluble supports for solution phase combinatorial chemistry (Padias, A.B.; Hall, H.K. Jr.; Tomalia, D.A.; McConnell, J.R. *J. Org. Chem.*, 52, 5305-5312 (1987); Gitsov, I.; Fréchet, J. M. J. *J. Am. Chem. Soc.*, 118, 3785-3786 (1996)).

The use of a soluble polyvalent support in accordance with the present invention provides significant advantages with respect to solid phase synthesis of combinatorial libraries, including: (1) solution phase synthesis obviates the need to modify chemistry to the solid phase; (2) intermediates may be routinely characterized by a variety of analytical methods, including  $^1\text{H}$  and  $^{13}\text{C}$  NMR, IR, UV and mass spectrometry, with the generation of NMR spectra of generally high resolution being made possible; (3) because multiple copies of each molecule are synthesized per polyvalent support, extremely high loadings may be attained; (4) size-based purification is general, since it does not rely on other physical differences between support-bound compounds and reagents; the use of large reagent excesses are also permitted; and (5) polyvalent supports offer a flexible framework that may be engineered to exhibit properties necessary for their desired applications. Combinatorial

chemistry on soluble polyvalent supports thus provides a valuable alternative to solid phase synthesis.

Commercially available Starburst™ polyamidoamine (PAMAM) dendrimers have been employed as a soluble support for combinatorial synthesis. However, PAMAM is not an ideal support for solution-phase chemistry, since the amine and amide moieties may be reactive under some conditions, and variable solubility of the dendrimers in organic solvents may limit the scope of chemistry which can be performed on the PAMAM supports. In addition, broad <sup>1</sup>H NMR signals observed for the PAMAM species could hinder characterization of bound intermediates.

Accordingly, the present compounds provide several advantages over previously used PAMAM dendrimers which are important for their use as soluble supports for combinatorial chemistry, including: (1) greatly enhanced solubility in a range of solvents, which broadens the scope of chemistry that can be performed, (2) significantly sharper <sup>1</sup>H NMR signals, which facilitates characterization of bound intermediates, and (3) higher chemical stability, which expands the scope of chemistry that can be carried out with such polyvalent supports.

## SUMMARY OF THE INVENTION


The present invention is directed to certain polyaryl-poly(ethylene glycol) compounds which are useful as soluble polyvalent supports in the preparation of combinatorial libraries of compounds. These polyaryl-PEG compounds are comprised of polyaryl cores bearing poly(ethylene glycol) chains which are attached through alkyl spacers. Synthons can be attached directly to the poly(ethylene glycol) termini, or to chemical linkers covalently attached to the poly(ethylene glycol) chains. These soluble polyvalent supports are useful in the preparation of a combinatorial library by facilitating the use of standard solution phase chemistry, and allowing homogeneous purification, routine characterization of intermediates, and high support loading.

The present invention is further directed to a combinatorial library of compounds wherein the combinatorial library contains such

compounds as a soluble polyvalent support. The present invention further embraces a combinatorial library of compounds which comprises a plurality of compounds attached to these soluble polyvalent supports.


## 5 DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to a compound of the Formula I, II or III:



wherein:

- 10 A, A', A'' and A''' are independently selected from:



X and X' are independently selected from the group consisting of:

-OH, -Br, -Cl, -I, -SH, -CO<sub>2</sub>H, -CO<sub>2</sub>Cl, CO<sub>2</sub>Br, -SO<sub>3</sub>H,  
-SO<sub>2</sub>Cl, -NH<sub>2</sub>, and -CH=CH<sub>2</sub>;

15

Y is independently selected from the group consisting of:

hydrogen, -CH<sub>3</sub>, -O-(C<sub>1-6</sub>alkyl) and -O(CH<sub>2</sub>CH<sub>2</sub>O)<sub>n</sub>-CH<sub>3</sub>;

j and j' are independently an integer from 0 to 4, inclusive;

20

k and k' are independently an integer from 1 to 12, inclusive;

m and m' are independently an integer from 1 to 4, inclusive; and

n is an integer from 0 to 12, inclusive;

and salts thereof.

5 Preferred compounds of the instant invention include those of Formula I, II and III wherein j and j' are 1, 2 or 3. Even more preferred compounds of the instant invention include those of Formula I, II and III wherein j and j' are 1 or 3.

10 Preferred compounds of the instant invention include those of Formula I, II and III wherein k and k' are 3 to 7, inclusive. Even more preferred compounds of the instant invention include those of Formula I, II and III wherein k and k' are 3, 4 or 5.

15 Preferred compounds of the instant invention include those of Formula I, II and III wherein m and m' are 2 or 3. Even more preferred compounds of the instant invention include those of Formula I, II and III wherein m and m' are 2.

Preferred compounds of the instant invention include those of Formula I, II and III wherein Y is independently selected from the group consisting of: hydrogen, -CH<sub>3</sub>, and -OCH<sub>3</sub>.




20 Preferred compounds of the instant invention include those of Formula I, II and III wherein X and X' are independently selected from the group consisting of: -OH, -Br, -NH<sub>2</sub>, and -CO<sub>2</sub>H. Even more preferred compounds of the instant invention include those of Formula I, II and III wherein X and X' are -OH.


25 Preferred compounds of the instant invention include those of Formula I, II and III wherein Y is independently selected from the group consisting of: hydrogen, and -OCH<sub>3</sub>.

30 In the compounds of Formula I, II and III, it is preferred that A, A', A'' and A''' (if present in the compound of Formula I, II and III) are identical to each other, i.e. it is preferred that the compounds are symmetrical.



Preferred compounds of the instant invention include:





As will be appreciated by those skilled in the art, the compounds of Formula I are biphenyl compounds, the compounds of Formula II are 1,3,5-triphenylbenzene compounds, and the compounds of Formula III are 3,5,3',5'-tetraphenylbiphenyl compounds.

The present invention envisions the use further modification of the present soluble polyvalent supports to facilitate manipulation and/or identification. Accordingly, the present invention is further directed to compounds wherein the compound of Formula I, II or III is covalently attached to a handle or linker functionality which facilitates the manipulation of the support or which acts as an anchor to facilitate the attachment/detachment of compounds thereto. Such handles or linkers include those which are known for use in solid phase synthesis, including 2,4-dimethoxy-4'-hydroxy-benzophenone, 4-(4-hydroxymethyl-3-methoxyphenoxy)-butyric acid (HMPBA), 4-hydroxymethylbenzoic acid,

4-hydroxymethyl-phenoxyacetic acid, 3-(4-hydroxymethylphenoxy)-propionic acid, p-[(R,S)- $\alpha$ -[1-(9H-fluoren-9-yl)methoxyformamido]-2,4-dimethoxy-benzyl]-phenoxyacetic acid, p-chloromethylphenyl linker, p-hydroxymethylphenyl linker, MBHA linker, HMBA-MBHA linker,  
5 Wang linker, Nova Syn TGA linker, Rink acid linker, Rink amide linker, Rink amide MBHA linker, Sieber liker, trityl linker, and the like.

In addition, the present invention is further directed to compounds wherein the compound of Formula I, II or III is attached to functionality which acts as a label or tag to facilitate the identification or  
10 characterization of the support (and any compounds attached thereto). Such labels or tags include chemical or radiochemical moieties which are capable of being distinguished from other chemical moieties and which are capable of being detected at low levels (such as at  $10^{-18}$  to  $10^{-19}$  mole), such as fluorescent labels. Representative tags include those  
15 disclosed in PCT Patent Publications WO 95/16209, WO 95/28640, WO 97/14814 and WO 97/15390.

The present invention is also directed to a method for the use of the instant compounds in the preparation of a combinatorial library.

The present invention is further directed to a combinatorial  
20 library wherein the combinatorial library is prepared by employing a compound of Formula I, II or III.

Accordingly, the present invention further embraces a combinatorial library which comprises a plurality of compounds attached to a compound of Formula I, II or III.

25 The present invention further embraces a combinatorial library wherein the combinatorial library is prepared by employing a compound of Formula I, II or III.

The present invention is further directed to a kit for combinatorial synthesis which comprises a compound of Formula I, II or  
30 III.

In an embodiment of the present invention the instant polyaryl-poly(ethylene glycol) soluble supports may be used for the preparation of combinatorial libraries. In a preferred embodiment the present invention is directed to a process for the preparation of

combinatorial libraries which comprises: (1) attaching a first synthon to a soluble polyvalent support to form a support-bound synthon; (2) separating the support-bound synthon from the reaction media; (3) optionally, mixing the support-bound synthon with one or more different support-bound synthons; (4) reacting the support-bound synthon or mixture of support-bound synthons with another synthon to form a support-bound product or mixture of support-bound products; (5) separating the support-bound product or mixture of support-bound products from the reaction media. In the above process, the order of steps (2) and (3) may be interchanged. In the above process, steps (3)-(5) may be repeated multiple times to afford the desired product or products. Optionally, the support-bound product or mixture of support-bound products may be cleaved and isolated from the support.

In a preferred embodiment, the present invention is directed to such a process which is conducted by automation or under computer control, since both the synthesis and purification steps are performed in solution.

By the term "polyvalent support" is meant a compound of Formula I, II or III.

By the term "soluble" is meant that the polyvalent support is dissolved in the appropriate solvent selected for the reaction media or subsequent manipulations. If desired, the polyvalent support may be soluble in the reaction media and insoluble in the solvents employed in a subsequent manipulation to facilitate isolation of the support by e.g. precipitation.

By the term "synthon" is meant any chemical moiety which may be synthetically manipulated to permit its covalent linking to a support or to another synthon. To facilitate the separation of the synthon from the polyvalent support it is preferred that the synthon be attached via a chemically cleavable linker. Upon cleavage from the support, the linked synthons comprise discrete molecular entities which may be analyzed for their biological activity or physiochemical properties, or which may be subjected to further chemical modification.

The compounds of the present invention are capable of forming salts with various inorganic and organic acids and bases and such salts are also within the scope of this invention. Examples of such acid addition salts include acetate, adipate, benzoate, benzenesulfonate, 5 bisulfate, butyrate, citrate, camphorate, camphorsulfonate, ethanesulfonate, fumarate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, methanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, oxalate, pamoate, persulfate, picrate, pivalate, propionate, succinate, tartrate, tosylate, and 10 undecanoate. Base salts include ammonium salts, alkali metal salts such as sodium, lithium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine and so forth. Also, the basic nitrogen- 15 containing groups may be quaternized with such agents as: lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl; diamyl sulfates; long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; aralkyl halides like benzyl bromide and others. 20 Such salts are useful for convenience in synthesis or for in isolating or purifying the product.

The salts may be formed by conventional means, such as by reacting the free base form of the compound with one or more equivalents of the appropriate acid in a solvent or medium in which the salt is 25 insoluble, or in a solvent such as water which is removed *in vacuo* or by freeze drying or by exchanging the anions of an existing salt for another anion on a suitable ion exchange resin.

In the present invention, the polyvalent support may be separated from solvents and reagents or otherwise purified by methods 30 known in the art including, crystallization, precipitation, ultrafiltration, gel permeation chromatography (GPC) (i.e. size exclusion chromatography (SEC)), precipitation, centrifugation, dialysis or selective adsorption. Preferred methods include ultrafiltration, size exclusion chromatography, precipitation, centrifugation and selective

adsorption, in which the most preferred methods include ultrafiltration, size-exclusion chromatography and precipitation.

The extremely high loadings capable on a polyvalent support greatly facilitates the production of multimilligram quantities of compounds. The ability to achieve high support loading increases efficiency by diminishing the physical size of the reactor which is required for the preparation of useful amounts of the desired compounds.

A combinatorial library is a collection of compounds in which the compounds comprising the collection are composed of one or more subunits or monomeric units (i.e. synthons). The subunits may be selected from natural or unnatural moieties including amino acids, nucleotides, sugars, lipids, carbohydrates, dienes, dienophiles, and the like. The compounds of the combinatorial library differ in one or more ways with respect to the type(s), number, order or modification of the subunits comprising the compounds.

The combinatorial libraries generated by the methods of the present invention may be screened for pharmacologically or diagnostically useful compounds, as well as for desired physical or chemical properties. It will be clear to one skilled in the art that such screening may be conducted on a library of compounds which have been separated from the polyvalent support, or may be conducted directly on the library of compounds which are still linked to the polyvalent support.

The present invention provides a general method by which any variety of single compounds or libraries may be created. Furthermore, polyvalent supports present a modular framework which can be custom-tailored so as to be applicable to a variety of synthetic methodologies.

The present invention is useful for developing new drugs and chemical entities. The invention is useful for rapidly generating and systematically synthesizing large numbers of molecules that may vary in their chemical structure or composition. The invention is further useful for randomly generating a large number of candidate compounds, then later optimizing those compounds which exhibit the most desirable properties.

As will be readily apparent to one skilled in the art, the present invention has application in essentially any synthetic reaction which may be conducted in the solution phase. Thus, the present invention is useful in almost all of the synthetic reactions which are

5 known to one of skill in the art, including peptide synthesis, acylation, alkylation, condensation, cyclization, halogenation, heterogeneous catalysis, hydrolysis, metallation, nitration, nucleophilic displacement, organometallic reactions, oxidation, reduction, sulfonation, acid chloride formation, Diels-Alder reaction, Friedel-Crafts reactions, Fischer indole

10 synthesis, Michael reactions, and the like (see e.g., H. O. House, "Modern Synthetic Reactions", 2nd ed. (Benjamin/Cummings, Menlo Park 1972); J. March, "Advanced Organic Chemistry", 3rd ed., (John Wiley & Sons, New York, 1985); Fieser and Fieser, "Reagents for Organic Synthesis", Volumes 1-end (Wiley Interscience, New York)). Likewise, the present

15 invention has application in essentially any synthetic reaction which may be conducted on solid phase supports, including acetal formation, alkylations, alkynylation, chiral alkylation, reductive alkylation, carbanion reactions, grignard reactions, organocadmium/manganese reactions, organolithium reactions, organozinc reactions, carbene insertion,

20 condensations, Claisen reactions, aldol reactions, Dieckmann cyclization, Knoevenagel condensations, mannich reactions, cycloadditions, cyclizations (in particular to form heterocyclic rings), Friedel-Crafts reactions, halogenation, bromination, chlorination, nucleophilic addition, Michael addition, aromatic nucleophilic substitution, Finkelstein reaction,

25 Mitsunobu reaction, palladium (0) catalyzed reactions, Stille coupling, Suzuki coupling, Heck reaction, carbamate/urea formation, oxidation of primary alcohol to aldehyde, Sharpless reaction, oxidation of secondary alcohol to ketone, oxidation of aldehyde to carboxylic acid, epoxidation, oxidation of primary chloride to aldehyde, oxidative phenol coupling,

30 reduction of acid to alcohol, reduction of aldehyde to alcohol, reduction of alkyne to alkene, reduction of amide to amine, reduction of aryl nitro to amine, reduction of azide to amine, reduction of ester to alcohol, reduction of imine to amine, reduction of iodide to alkyl, reduction of ketone to alcohol, Wittig reaction, Horner-Emmons condensation, and the


like (see generally, "Solid Phase Organic Chemistry (SPOC)" and "Solid Phase Inorganic Chemistry (SPIC)", Chiron Mimotopes, pp. 1-31 (August 1995).

As outlined schematically in Scheme 1, the present  
5 compounds may be employed as a soluble polyvalent support  
analogously to a solid phase support in traditional combinatorial  
synthesis, except that reactions are performed in solution, and  
intermediates bound to the polyvalent supports are separated by methods  
such as ultrafiltration, size exclusion chromatography, precipitation,  
10 centrifugation, dialysis or selective adsorption, and especially size-  
selective methods such as gel permeation chromatography or  
ultrafiltration. As will be clear to one skilled in the art, the steps may be  
repeated iteratively to incorporate additional synthons ( $R^3$ ,  $R^4$ , etc) into  
the target compound (i.e.  $R^1$ - $R^2$ - $R^3$ - $R^4$ , ... etc.).


15



**SCHEME 1**



As used above, the generic depiction of a polyvalent support



is not intended to represent any specific arrangement of carbon atoms, but is merely a graphical representation of the compound of Formula I, II or III which representation is employed for convenience and to facilitate ready comprehension of the essential features of the present invention.

The preparation of compounds of Formula I, II and III of the present invention may be carried out in sequential or convergent synthetic routes. Syntheses detailing the preparation of the compounds of Formula I, II and III in a sequential manner are presented in the following reaction schemes.

The general preparation of compounds of the present invention is described in the following Reaction Schemes.


Polyaryl-PEG supports may be prepared by either convergent or divergent schemes. A general convergent approach toward polyaryl-PEG production involves: (1) attachment of unprotected or monoprotected PEG chains to an aryl bromide or iodide, (2) conversion of the aryl halide to an aryl species capable of undergoing cross-coupling to an aryl halide (i.e. an aryl stannane or aryl boronic acid), (3) cross coupling of the aryl species with an aryl halide to form the polyaryl-PEG compound, and, if necessary, (4) deprotection of the PEG chains to afford the polyaryl-PEG support.

Specific examples of convergent schemes are detailed in the following examples. Alternatively, divergent schemes involving attachment of PEG chains to preformed polyaryl cores may also be used to prepared the instant polyaryl-PEG supports.

Asymmetric polyaryl-PEG supports may also be prepared. The chains on a given A may be differentiated by stepwise coupling of the aryl group with a single PEG chain (k, m, X), followed by coupling of


- a second chain ( $k'$ ,  $m'$ ,  $X'$ ). X-groups may also be differentiated after attachment of the PEG chains to the aryl moiety by conversion of one of the X groups to  $X'$  (with employment of protecting groups as required). Polyaryl-PEG supports composed of different A-groups may also be prepared. One approach is to couple asymmetric aryl compounds to afford the asymmetric polyaryl species. For example, coupling of  $A-SnMe_3$  with  $A'-Br$  yields asymmetric biphenyl  $A-A'$ , while stepwise coupling of  $A-SnMe_3$ ,  $A'-SnMe_3$  and  $A''-SnMe_3$  with 1,3,5-tribromobenzene yields the asymmetric triarylbenzene. Differences in reactivity of aryl species toward cross-coupling reactions (for example, the greater propensity of aryl iodides to undergo Stille cross-coupling reactions vs. aryl bromides) may also be exploited in the stepwise production of asymmetric polyaryl-PEG supports of types II and III. Asymmetry may also be conferred to the polyaryl-PEG supports by introduction of different PEG chains to a polyaryl core. This process would be facilitated by differentiating sites of PEG attachment on the polyaryl core (i.e. by employing benzyl bromide vs. benzyl chloride groups).

SCHEME 2




- As outlined in Scheme 2, a polyethylene glycol bearing the desired functionality at X may be protected with an appropriate protecting group (if desired), then coupled with 3,5-bis-bromomethylbromobenzene to give the 3,5-bis(PEG)bromobenzene compound. If unsymmetrical compounds are desired (i.e. of the formula j', k', m' and X'), 3-bromomethyl-5-iodomethylbromobenzene may be reacted first with one polyethylene glycol, followed by reaction with a different polyethylene glycol. Due to the sensitive nature of the functionality at X, the use of protecting groups will generally be necessary. Alternatively, the desired functionality at X may be introduced at a later stage by synthetic modification of an appropriate polyaryl precursor.

**SCHEME 3**




As outlined in Scheme 3, the aryl bromide is converted to the aryl stannane using hexamethylditin and catalytic tetrakis(triphenylphosphine)palladium(0) in refluxing toluene.

**SCHEME 4**




5 As outlined in Scheme 4, Pd-catalyzed Stille cross-coupling (or alternatively Suzuki coupling) of the aryl stannane with its precursor aryl bromide (and removal of protecting groups if present) to give the desired biphenyl compound. As will be readily apparent to one skilled in the art, the biphenyl compound may be symmetrical or unsymmetrical. Unsymmetrical biphenyl compounds may be prepared by reacting the aryl stannane with an aryl bromide which bears different substituents.

**SCHEME 5**



- As outlined in Scheme 5, Pd-catalyzed Stille cross-coupling of the aryl stannane with 1,3,5-tri-bromobenzene (and removal of protecting groups if present) gives the desired 1,3,5-triaryl benzene compound. As will be readily apparent to one skilled in the art, the 1,3,5-triphenylbenzene compound may be symmetrical or unsymmetrical (i.e. bearing A, A', A''). Unsymmetrical 1,3,5-triphenylbenzene compounds may be prepared by reacting the phenyl tin derivative with a halo-benzene (e.g. 1-iodo-3,5-dibromobenzene) followed by reaction with a different phenyl tin derivative.

**SCHEME 6**



As outlined in Scheme 6, Pd-catalyzed Stille cross-coupling of the aryl stannane with 3,5,3',5'-tetrabromobiphenyl (and removal of protecting groups if present) gives the desired 3,5,3',5'-tetraaryl biphenyl compound. As will be readily apparent to one skilled in the art, the 3,5,3',5'-tetraphenyl biphenyl compound may be symmetrical or unsymmetrical (i.e. bearing A, A', A'', A''').



5

HO(CH2CH2O)3CH2CH2OH + C1=CC=CC=C1O1 >> HO(CH2CH2O)3CH2CH2OC1=CC=CC=C1O1


TsOH/CH<sub>2</sub>Cl<sub>2</sub>


**1**

(1) NaH/THF  
 (2) BrCc1ccc(Br)cc1

BrCc1ccc(Br)cc1

**2**






As depicted in Scheme 7, a convergent approach was used to prepare polyaryl-PEG supports of Formula I and II where  $j = 1$ ,  $k = 3$ ,  $m = 2$  and  $X = -\text{OH}$  (or  $-\text{OTHP}$ ). Mono-THP protection of tetraethylene glycol, conversion to the sodium derivative and coupling with 3,5-bis-bromomethylbromobenzene afforded aryl bromide 2 bearing two PEG chains. Aryl bromide 2 was converted to the aryl stannane 3 using hexamethylditin and catalytic tetrakis(triphenylphosphine)palladium(0) in refluxing toluene. Pd-catalyzed Stille cross-coupling of 3 with either its precursor aryl bromide 2 or with 1,3,5-tribromobenzene or with 3,5,3',5'-

5

**SCHEME 8**



As depicted in Scheme 8, a sequential approach is used to prepare polyaryl-PEG supports of Formula I where  $j = 3$ ,  $k = 5$ ,  $m = 2$ ,  $Y = -OCH_3$  and  $X = -OH$  (or  $-OTHP$ ). Alkylation of 4,4'-biphenol with allylbromide in DMF containing  $Cs_2CO_3$  afforded allyl ether 7, which  
5 was converted to 3,3'-bis-allyl-4,4'-biphenol 8 by heating neat at  $190^\circ C$  for 18 h. under nitrogen. Analogous re-allylation of biphenol 8, followed by Claisen rearrangement, yielded tetraallyl compound 10. Alkylation of 10 with MeI in presence of  $K_2CO_3$  in DMF at  $40^\circ C$  yielded dimethoxy species 11. Treatment of the tetraolefin with 9-BBN in THF, followed by  
10 hydrogen peroxide oxidation afforded tetra-ol 12, which was converted to tetrabromide 13 using NBS and  $PPh_3$  in  $CH_2Cl_2$ . Formation of protected support 14 was achieved by nucleophilic displacement of the tetrabromide with mono-THP protected hexa(ethylene glycol) using NaOH in THF in presence of a phase-transfer catalyst. The deprotection  
15 of the THP-protected compounds is effected with MeOH containing catalytic toluenesulfonic acid to afford the corresponding alcohol.

Many of the starting materials are either commercially available or known in the literature and others can be prepared following literature methods described for analogous compounds. The skills  
20 required in carrying out the reaction and purification of the resulting reaction products are known to those in the art. Purification procedures include crystallization, ultrafiltration, size exclusion chromatography (gel permeation chromatography), precipitation, centrifugation, dialysis, selective adsorption, chromatography on silica gel (including flash  
25 chromatography), and normal phase or reverse phase chromatography.


As will be appreciated by one skilled in the art, protecting groups may be employed as required. The term "protecting group" is intended to indicate the presence of an appropriate protecting group, such as those described in Greene, T.W., Wuts, P.G.M. "Protective Groups in  
30 Organic Synthesis", 2nd ed., John Wiley & Sons, Inc., New York, 1991. An appropriate protecting group will be able to withstand the reaction conditions of intermediate processes, prior to being removed when desired. Suitable protecting groups for hydroxyl include those groups well known in the art such as: tetrahydropyranyl; 1-(lower

- alkylthio)(lower)alkyl, wherein "lower alkyl" indicates a straight, cyclic or branched chain of one to six carbon atoms, such as lower alkylthiomethyl (e.g. methylthiomethyl, ethylthiomethyl, propylthiomethyl, isopropylthiomethyl, butylthiomethyl, isobutylthiomethyl, hexylthiomethyl, etc.), and the like; trisubstituted silyl such as tri(lower)alkylsilyl (e.g. trimethylsilyl, triethylsilyl, tributylsilyl, tri-i-propylsilyl, t-butyl dimethylsilyl, tri-t-butylsilyl, etc.), lower alkyl diarylsilyl (e.g. methyl-diphenylsilyl, ethyl-diphenylsilyl, propyl-diphenylsilyl, t-butyl diphenylsilyl, etc.), and the like; acyl such as
- 5 aliphatic acyl, aromatic acyl and aliphatic acyl substituted with aromatic group, which are derived from carboxylic acids; and the like. Suitable protecting groups for amino include those groups well known in the art such as: benzyl, benzyloxymethyl, benzyloxycarbonyl (carbobenzyloxy), benzylsulfonyl, 2-bromoethyloxycarbonyl, t-butoxy-carbonyl, 2-chloro-
- 10 benzyloxycarbonyl, 2-chloroethyloxycarbonyl, di-t-amylloxycarbonyl, 9-fluoroenylmethyloxycarbonyl, isopropoxycarbonyl, 4-methoxy-benzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 2-nitrophenyl-sulfonyl, phthaloyl, 2,2,2-trichloro-t-butyloxycarbonyl, trifluoroacetyl, tirphenylmethane, and vinyloxycarbonyl groups, and the like.
- 15
- 20 Methodology for the introduction and removal of protecting groups is well known in the art (see e.g. Greene, T.W., Wuts, P.G.M. "Protective Groups in Organic Synthesis", 2nd ed., John Wiley & Sons, Inc., New York, 1991).

- The following examples are provided for the purpose of
- 25 further illustration only and are not intended to be limitations on the disclosed invention.



**EXAMPLE 16**



**3,3',5,5'-Tetra-[hexa(ethyleneglyoxy)]-4,4'-dimethoxybiphenyl ether (15)**


- The tetra-THP 14 is deprotected essentially as described in  
5 Example 5 with a 0.02 M toluenesulfonic acid solution in MeOH at room temperature to give the title compound.

While the invention has been described and illustrated with reference to certain particular embodiments thereof, those skilled in the art will appreciate that various adaptations, changes, modifications, substitutions, deletions, or additions of procedures and protocols may be made without departing from the spirit and scope of the invention. For example, reaction conditions other than the particular conditions as set forth herein above may be applicable as a consequence of variations in the reagents or methodology to prepare the compounds from the processes of the invention indicated above. Likewise, the specific reactivity of starting materials may vary according to and depending upon the particular substituents present or the conditions of manufacture, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. It is intended, therefore, that the invention be defined by the scope of the claims which follow and that such claims be interpreted as broadly as is reasonable.




WHAT IS CLAIMED IS:

1. A compound of the Formula I, II or III:



5 wherein:

A, A', A'' and A''' are independently selected from:



X and X' are independently selected from the group consisting of:

- 10                      - OH, -Br, -Cl, -I, -SH, -CO<sub>2</sub>H, -CO<sub>2</sub>Cl, CO<sub>2</sub>Br, -SO<sub>3</sub>H,  
                         -SO<sub>2</sub>Cl, -NH<sub>2</sub>, and -CH=CH<sub>2</sub>;

Y is independently selected from the group consisting of:

hydrogen, -CH<sub>3</sub>, -O-(C<sub>1</sub>-6alkyl) and -O(CH<sub>2</sub>CH<sub>2</sub>O)<sub>n</sub>-CH<sub>3</sub>;

- 15    j and j' are independently an integer from 0 to 4, inclusive;  
      k and k' are independently an integer from 1 to 12, inclusive;  
      m and m' are independently an integer from 1 to 4, inclusive; and  
      n is an integer from 0 to 12, inclusive;

20    and salts thereof.

2. The compound of Claim 1 wherein j and j' are  
1, 2 or 3.
3. The compound of Claim 2 wherein j and j' are  
1 or 3.
4. The compound of Claim 1 wherein k and k' are  
3 to 7, inclusive.
5. The compound of Claim 4 wherein k and k' are  
3, 4 or 5.
6. The compound of Claim 1 wherein m and m' are  
2 or 3.
7. The compound of Claim 6 wherein m and m' are 2.
8. The compound of Claim 1 wherein Y is independently  
selected from the group consisting of: hydrogen, -CH<sub>3</sub>, and -OCH<sub>3</sub>.
9. The compound of Claim 1 wherein X and X' are  
independently selected from the group consisting of: - OH, -Br, -NH<sub>2</sub>,  
and -CO<sub>2</sub>H.
10. The compound of Claim 1 wherein Y is independently  
selected from the group consisting of: hydrogen, and -OCH<sub>3</sub>.
11. The compound of Claim 1 wherein X and X' are  
- OH.
12. The compound of Claim 1 wherein j and j' are  
1 or 3, k and k' are 3, 4 or 5, and m and m' are 2.

13. The compound of Claim 12 wherein Y is independently selected from the group consisting of: hydrogen, and -OCH<sub>3</sub>.

5 14. The compound of Claim 12 wherein X and X' are - OH.


10 15. The compound of Claim 1 wherein A, A', A" and A''' (if present in the compound of Claim 1) are identical to each other.


16. The compound of Claim 1 which additionally comprises a linker functionality which is selected from the group consisting of: 2,4-dimethoxy-4'-hydroxy-benzophenone, 4-(4-hydroxymethyl-3-methoxyphenoxy)-butyric acid, 4-hydroxymethylbenzoic acid, 4-hydroxymethyl-phenoxyacetic acid, 3-(4-hydroxymethylphenoxy)-propionic acid, p-[(R,S)- $\alpha$ -[1-(9H-fluoren-9-yl)methoxyformamido]-2,4-dimethoxy-benzyl]-phenoxyacetic acid, p-chloromethylphenyl linker, p-hydroxymethylphenyl linker, MBHA linker, HMBA-MBHA linker, Wang linker, Nova Syn TGA linker, Rink acid linker, Rink amide linker, Rink amide MBHA linker, Sieber liker, and trityl linker.

15


20

5





and



5

18. A method for using the compound of Claim 1 in the preparation of a combinatorial library.

10

19. A combinatorial library which comprises a plurality of compounds attached to a compound of Claim 1.

20. A kit for combinatorial synthesis which comprises a compound of Claim 1.

